

UCT-TP 191/93
27 January 1993

Indications for a Detonating Quark-Gluon Plasma

N. Bilić^{1,2}, J. Cleymans¹, E. Suhonen^{1,3}, and D.W. von Oertzen⁴

¹Department of Physics, University of Cape Town, Rondebosch 7700, South Africa

²Rudjer Bošković Institute, P.O. Box 1016, Zagreb, Croatia

³Department of Theoretical Physics, University of Oulu, SF-90550 Oulu, Finland

⁴Physics Department, University of Namibia, Windhoek, Namibia

Abstract

We propose a mechanism which naturally contains the relation $\mu_B = 3\mu_S$ of the hadronic gas produced in heavy-ion collisions at CERN. Our starting assumption is the existence of a sharp front separating the quark-gluon plasma phase from the hadronic phase. Energy-momentum conservation across the front leads to the following consequences for an adiabatic process

- a) The baryon chemical potential, μ_B , is approximately continuous across the front.
- b) The temperature in the hadronic gas is higher than the phase transition temperature due to superheating.
- c) In the region covered by the experiments the velocity of the hadronic gas approximately equals the speed of sound in the hadronic gas.

Several groups [1]-[8] have investigated the results obtained by the CERN relativistic ion experiments on the production of strange particles [9]-[12]. These results are very sensitive to the particular kinematic window in which particle yields are being measured. The results from the WA85 group [10] have attracted particular attention because they measure several of the strange baryons in a very narrow kinematic region, central in rapidity and at large transverse momenta. Within the framework of hadronic gas models these experimental results lead to the conclusion that the hadrons were produced from a system having a temperature T of approximately 200 MeV, a baryon chemical potential μ_B of about 300 MeV and a strangeness chemical potential μ_S of 100 MeV [1, 3, 8]. The ratio of strange to baryon chemical potential thus obtained is particularly intriguing as the relation

$$\mu_B = 3\mu_S \tag{1}$$

naturally holds in a quark-gluon plasma [1] guaranteeing strangeness neutrality between strange and anti-strange quarks. In a hadronic gas there is however no particular reason for such a relation to hold. Early proposals to explain it have difficulties with the entropy generated in the transition and violate the second law of thermodynamics [2].

In the present paper we propose a model which is obtained naturally if one assumes the existence of a sharp front separating the hadronic and quark-gluon plasma phases. Ahead and behind the front each phase is behaving as an (approximately) free gas of particles, except for the fact that we take into account the hard core of the baryons and the presence of resonances. All the dynamics related to the transition takes place inside this sharp front. In going from one phase to the other we take into account energy-momentum conservation. This provides for a certain relationship between the energy-density in the hadronic phase and in the quark-gluon phase. In order for the transition to be possible, a substantial amount of supercooling has to happen in the quark-gluon phase while a super-heated hadronic gas emerges on the other side. This way there is no problem with the entropy since the supercooled quark-gluon gas will have a low entropy content. One is then led to the picture which is shown in figure 1. Such a transition is well known in hydrodynamics and is usually referred to as a detonation transition (see e.g. Ref. [13]). The idea of a sharp front was first put forward in this context by Van Hove [14], it was considered in more detail later on by several groups [15]-[20]. It was also considered recently in the context of particle production in Ref. [21]. It has however not been used in any systematic way to make contact with the recent experimental results obtained with the CERN relativistic ion experiments. This paper is a first step towards such an analysis.

The attractive features of our model are as follows :

1. The baryon chemical potential μ_B is continuous across the front to a very good approximation.
2. The relation $\mu_B = 3\mu_S$ is exactly preserved during the transition.
3. The temperature in the hadronic gas is higher than the phase transition temperature due to the resulting superheating. This would explain why numerical

simulations [22]-[24] of lattice QCD obtain estimates for the critical temperature which are lower than the temperature observed in the transverse momenta spectra measured in relativistic ion collisions.

4. The hadronic gas emerges approximately with the velocity of sound.

Imposing the condition $\mu_B = 3\mu_S$ in the hadronic phase constrains the allowed values of the temperature. Normally μ_S is a free parameter whose value is fixed by the condition of strangeness neutrality in the hadronic gas. Fixing it to the above relation implies imposing a restriction on the temperature. In fact, the hadronic gas cannot overheat beyond a certain limit (about 205 MeV in our case). The resulting curve in the $T - \mu_B$ plane is shown in figure 2. Note also that for $\mu_B = 0$ no constraint exists since μ_S is then also zero and the relation is trivially satisfied.

As stated before we assume the existence of a sharp front separating the two phases. The standard analysis considers a one-dimensional situation in the frame of reference where the front is at rest (see e.g. Ref. [13]). Energy and momentum conservation across the front then lead to the two following equations :

$$(\epsilon_h + P_h)\gamma_h^2 v_h + P_h = (\epsilon_q + P_q)\gamma_q^2 v_q + P_q, \quad (2)$$

$$(\epsilon_h + P_h)v_h^2\gamma_h^2 = (\epsilon_q + P_q)v_q^2\gamma_q^2, \quad (3)$$

both sides of the front are at different temperatures and chemical potentials. The thermodynamic quantities (pressure, energy density) refer to the rest frame quantities. The index q refers to the quark-gluon plasma phase while the index h refers to the hadronic phase, v refers to the velocity of the gas in each phase with respect to the front, γ is the standard relativistic factor ($1/\sqrt{1-v^2}$). In addition to the above two equations, baryon number is also conserved across the front which leads to

$$n_h v_h \gamma_h = n_q v_q \gamma_q. \quad (4)$$

where n_h and n_q refer to the baryon densities in the hadronic and in the quark-gluon plasma phases respectively. The above equations can be solved for the velocities, leading to

$$v_q^2 = \frac{(P_h - P_q)(\epsilon_h + P_q)}{(\epsilon_h - \epsilon_q)(\epsilon_q + P_h)}, \quad (5)$$

$$v_h^2 = \frac{(P_h - P_q)(\epsilon_q + P_h)}{(\epsilon_h - \epsilon_q)(\epsilon_h + P_q)}. \quad (6)$$

The change in density can then be determined from

$$\left(\frac{n_q}{n_h}\right)^2 = \frac{(\epsilon_q + P_q)(\epsilon_q + P_h)}{(\epsilon_h + P_h)(\epsilon_h + P_q)}. \quad (7)$$

Another important constraint is given by the requirement of non-decreasing entropy which implies in our notation

$$s_h v_h \gamma_h \geq s_q v_q \gamma_q. \quad (8)$$

Combining (4) and (8) leads to

$$\frac{s_h}{s_q} \geq \frac{n_h}{n_q}, \quad (9)$$

The equality sign is of particular interest since it corresponds to an adiabatic transition.

To exploit these relations fully we have to specify the equation of state (EOS) used in each phase. On the quark gluon plasma side we use the bag EOS given by

$$\epsilon_q = 3P_q + 4B, \quad (10)$$

where B is the bag constant. We have chosen $B = 0.2 \text{ GeV/fm}^3$.

On the hadronic side we incorporate the hard core radius of hadrons by choosing [25]

$$\epsilon_h = \frac{\epsilon_h^0}{1 + V_0 n_h^0}, \quad (11)$$

where the subscript zero refers to point-like quantities, V_0 refers to a typical proper volume of a hadron which we take to have a radius given by 0.8 fm . The hadronic side in our approach is a composition of non-interacting quantum gases. The use of quantum statistics is essential since in a wide range of μ_B and T the Boltzmann approximation is not satisfied. We include all well established hadronic resonances listed in the latest edition of the Review of Particle Properties [26]. A correction similar to equation (11) is performed to all thermodynamic quantities.

We will pay particular attention to adiabatic transitions corresponding to

$$\frac{s_q}{n_q} = \frac{s_h}{n_h}, \quad (12)$$

i.e. where the entropy per baryon remains constant across the transition. To determine the temperature in the quark phase we proceed as follows. Introducing the ratio z defined by

$$z = \mu_q/(\pi T_q), \quad (13)$$

one can use the equation of state described above for the quark-gluon phase to obtain

$$\frac{s_q}{n_q} = \frac{3\pi}{z} \left(1 - \frac{11}{180(1 + z^2)} \right). \quad (14)$$

We have chosen the number of flavours equal to 2.5 to simulate the effect of the strange quark mass while keeping the expressions for massless quarks [27]. This determines z for an adiabatic transition in terms of the hadronic temperature and baryon chemical potential. From the ratio of baryon densities given by equation (6) one can then determine the temperature T_q in the quark-gluon phase.

In figure 3 we show the energy densities in the quark and in the hadronic phases as functions of the temperature T_h . One sees that the energy density is always lower in the quark sector than in the hadronic sector for the range of temperatures considered.

In figure 4 we show the corresponding temperature in the quark phase. As one can see, it is always very low, indicating a substantial amount of supercooling in the quark phase.

Figure 5 is probably the most interesting result of this analysis. In this case we show the baryon chemical potential in the hadronic phase and the corresponding one on the quark side (denoted by $3\mu_Q$) varying the bag constant in the range 0.16-0.24 GeV/fm³. The two chemical potentials are nearly equal. This means that the baryon chemical potential is conserved through the front even though the temperatures are very different. So we have the interesting situation where all the thermodynamic quantities change quite drastically during the transition but the chemical potentials remain almost unaffected.

The velocities near the front have a special hydrodynamic importance. As is well known, small perturbations in a medium propagate with the velocity of sound in that medium. It is therefore of special interest to compare v_q and v_h with the velocity of sound in each phase. The latter is defined as [13]

$$c_s^2 = \left. \frac{\partial P}{\partial \epsilon} \right|_{s/n}. \quad (15)$$

In the quark-gluon phase, eq. (10) yields $c_{s,q}^2 = 1/3$. To calculate the speed of sound in the hadronic gas we write eq. (15) as

$$c_{s,h}^2 = \frac{\partial P_h / \partial T + \partial P_h / \partial \mu_B \cdot d\mu_B / dT}{\partial \epsilon_h / \partial T + \partial \epsilon_h / \partial \mu_B \cdot d\mu_B / dT}, \quad (16)$$

where the derivative $d\mu_B / dT$ is taken such that the ratio s_h / n_h remains constant. Thus

$$\frac{d\mu_B}{dT} = - \frac{n_h \partial s_h / \partial T - s_h \partial n_h / \partial T}{n_h \partial s_h / \partial \mu_B - s_h \partial n_h / \partial \mu_B}. \quad (17)$$

The velocities thus calculated are shown in figure 6 together with v_q and v_h . It can be seen that the quark-gluon plasma velocity is very high, quarks stream in at high speed but with a low density. On the other side of the front the hadrons are slower however with a larger density.

It is interesting to note that around the relevant values of T and μ_B namely 200 and 300 MeV respectively, the hadrons emerge with sound velocity, while the quarks have a velocity which is considerably larger than the speed of sound in a quark-gluon plasma.

One also observes in this model a large increase in the entropy density as one makes the transition from a supercooled quark-gluon phase to a superheated hadronic phase.

In summary, we have considered a model based on the existence of a sharp front separating the hadronic from the quark-gluon phase. We arrive at the conclusion that in this situation the transition can only take place if a substantial amount of super-cooling takes place in the quark-gluon phase and that the hadronic phase appears in a super-heated state. This would explain why such high temperature have been observed experimentally while the data from lattice QCD indicate that the phase transition temperature is considerably lower than 200 MeV. It is interesting to note that the chemical potentials remain approximately the same in such a transition while the temperature

changes drastically in the transition. The temperature on the hadronic side is about 100 MeV higher than on the quark-gluon side. In the frame where the sharp front is at rest the velocity of the quark phase is above the corresponding sound velocity while in the hadronic phase it is very close to the hadronic speed of sound.

Acknowledgement We acknowledge continuing stimulating discussions in this field with H. Satz and K. Redlich. Two of us (E.S. and D.W.v.O.) thank the Physics Department of the University of Cape Town for its hospitality.

References

- [1] J. Cleymans and H. Satz, Z. Phys. C, in press.
- [2] J. Cleymans, K. Redlich, H. Satz and E. Suhonen, CERN preprint CERN-TH 6759/02.
- [3] J. Rafelski, Phys. Lett. B262 (1991) 333.
- [4] J. Rafelski, Nucl. Phys. A544 (1992) 279c.
- [5] J. Letessier, J. Rafelski and A. Tounsi, Paris Preprint PAR-LPTHE 92-23, June 1992.
- [6] N.J. Davidson, H.G. Miller, R.M. Quick and J. Cleymans, Phys. Lett. B255 (1991) 195.
- [7] D.W. von Oertzen, N.J. Davidson, R.A. Ritchie and H.G. Miller, Phys. Lett. B274 (1992) 128.
- [8] N.J. Davidson, H.G. Miller, D.W. von Oertzen and K. Redlich, Z. Phys. C56 (1992) 319.
- [9] J. Bartke et al. (NA35), Z. Phys. C48 (1990) 191; R. Stock et al. (NA35), Nucl. Phys. A525 (1991) 221c.
- [10] S. Abatzis et al. (WA85), Phys. Lett. B244 (1990) 130; S. Abatzis et al. (WA85), Phys. Lett. B259 (1991) 508.
- [11] E. Andersen et al., Phys. Lett. B294 (1992) 127.
- [12] A. Franz (NA44), Report at the XXVI International Conference on High Energy Physics, 6.-12.8.1992, Dallas/Texas, USA.
- [13] L. Landau and E.M. Lifshitz, Hydrodynamics, Pergamon Press, London (1959).
- [14] L. Van Hove, Z. Phys. C21 (1983) 93; C27 (1985) 135.
- [15] M. Gyulassy, K. Kajantie, H. Kurki-Suonio and L. McLerran, Nucl. Phys. B237 (1984) 477.
- [16] J. Cleymans, E. Nykänen and E. Suhonen, Phys. Rev. D33 (1986) 2585.
- [17] J. Cleymans, E. Nykänen and E. Suhonen Z. Phys. C37 (1987) 51.
- [18] H.W. Barz, L.P. Csernai, B. Kämpfer and B. Lukács, Phys. Rev. D32 (1985) 115.
- [19] K.A. Bugaev, M. Gorenstein and V.I. Zhdanov, Z. Phys. C34 (1987) 365.
- [20] D. Seibert, Phys. Rev. D32 (1985) 2812; D35 (1987) 2013.

- [21] S.P. Baranov and L.V. Fil'kov, Z. f. Physik (to be published).
- [22] J.B. Kogut, Nucl. Phys. B270[FS16] (1986) 169.
- [23] F. Karsch, J.B. Kogut, D.K. Sinclair and H.W. Wyld, Phys. Lett. B188 (1987) 353.
- [24] N. Bilić, K. Demeterfi and B. Petersson, Nucl. Phys. B377 (1992) 651.
- [25] J. Cleymans, K. Redlich, H. Satz and E. Suhonen, Z. Phys. C33 (1986) 151.
- [26] Particle Data Group, Phys. Rev. D45 (1992) S1.
- [27] K. Kajantie and H. Kurki-Suonio, Phys. Rev. D34 (1986) 1719.

Figure Captions:

1. Schematic diagram of the energy-density as a function of temperature. A detonation transition from a supercooled quark-gluon plasma (long dashed line) to a superheated hadron gas (short dashed line) is indicated.
2. The hadronic temperature T_h as a function of the baryon chemical potential μ_B .
3. The energy-density of the hadronic gas (full line) and of the quark-gluon plasma (dashed line) as a function of the hadronic temperature.
4. The quark temperature T_q versus the hadronic temperature T_h .
5. The baryon chemical potential in the quark-gluon plasma for $B = 0.24 \text{ GeV/fm}^3$ (dash-dotted line), for $B = 0.2 \text{ GeV/fm}^3$ (full line), for $B = 0.16 \text{ GeV/fm}^3$ (long dashes) and in the hadronic gas (short dashes).
6. The velocity squared of the quark matter (long dashed line) and of the hadronic matter (short dashed line) relative to the transition front. Also shown are the hadronic sound velocity squared with (full line) and without (dot-dashed line) excluded volume corrections.